

# Rheumatoid Arthritis Treatment Guidelines

## 5-Star Best Practices



### Measure Description

The percentage of rheumatoid arthritis patients who were dispensed one or more prescriptions for an anti-rheumatic drug.

### Metric

The percentage of patients who were diagnosed with rheumatoid arthritis at least two times during the measurement year and who were dispensed at least one ambulatory prescription for a disease modifying anti-rheumatic drug (DMARD).

### How Physicians Can Improve on This 5-Star Measure

- > Confirm rheumatoid arthritis diagnosis versus a different rheumatologic disease, osteoarthritis or joint pain.
  - Best practice is to refer patients to a rheumatologist to confirm diagnosis and/or co-manage.
  - Consider using ICD-10 codes, such as M25.5x, for generalized joint pain while the patient's pain is being investigated. Use ICD-10 codes M05x and M06x once rheumatoid arthritis is confirmed. Other codes to consider when rheumatoid arthritis has not been established:
    - M19.0x – Other and unspecified osteoarthritis
    - M25.5x – Pain in joint
    - M25.6x – Stiffness of joint
    - M25.8x – Other specified joint disorders
    - M25.9x – Joint disorder, unspecified
  - To reinstate an encounter with an incorrect diagnosis code for rheumatoid arthritis, contact the medical group.
- > Evaluate the risks, benefits and alternatives of DMARD for all patients with rheumatoid arthritis and prescribe when appropriate.

Description	DMARD Prescription		
<b>5-Aminosalicylates</b>	Sulfasalazine (T2)		
<b>Alkylating agents</b>	Cyclophosphamide capsules (T2)		
<b>Aminoquinolines</b>	Hydroxychloroquine (T2)		
<b>Anti-rheumatics</b>	Auranofin (T5) Gold sodium thiomalate (NF)	Leflunomide (T2) Methotrexate solution (inj) (T2)	Methotrexate tablets (T2) Penicillamine (T4)
<b>Immunomodulators</b>	Abatacept (NF) Adalimumab (T5) Anakinra (T5) Certolizumab (NF)	Certolizumab pegol (NF) Etanercept (T5) Golimumab (NF) Infliximab (T5)	Rituximab (T5) Tocilizumab (NF)
<b>Immunosuppressive agents</b>	Azathioprine (T2) Cyclosporine (T2)	Mycophenolate suspension (T5)	Mycophenolate tablets/capsules (T2)
<b>Janus kinase (JAK) inhibitor</b>	Tofacitinib (T5)		
<b>Tetracyclines</b>	Minocycline immediate release (T2)		

Table above is a list of DMARDs in the SCAN Formulary. Tier 2 (T2) drugs are generic with lower copays. Tier 5 (T5) are specialty tier drugs with various coinsurance for a 30-day supply. Prescriptions marked with (NF) are not in SCAN's Formulary.

- See SCAN's Formulary for further information: <https://www.scanhealthplan.com/providers/>
- A sample DMARD dispensed in the office will satisfy the measure but it must be documented properly.
  - Record in the medical record that a DMARD was dispensed. **Important:** You must include all of the following: prescriber's signature, drug name, strength, days' supply and dispensed date.
- > For American College of Rheumatology guidelines, visit <https://www.rheumatology.org/practice-quality/clinical-support/criteria>.

# Rheumatoid Arthritis Medical Group Guidelines

## 5-Star Best Practices



### How the Managing Group Can Improve on This 5-Star Measure

- > Provide all your physicians with provider instructions located on the reverse side.
- > Best practice is to have a rheumatologist review all rheumatoid arthritis diagnoses.
- > Ensure there is adequate access to rheumatologists for patients to schedule appointments.
- > Document and distribute to your physicians what is needed to correct a rheumatoid arthritis diagnosis.
  - Correct any incorrect rheumatoid arthritis diagnosis.
    - Diagnosis codes can be corrected through SCAN's provider portal or by submitting an ICE file.
    - Ensure appropriate staff has access to the provider portal.
    - Contact your SCAN encounter team representative for instructions or questions on how to use the provider portal or how to submit an ICE file at [NetworkQuality@scanhealthplan.com](mailto:NetworkQuality@scanhealthplan.com).

Contact your SCAN network management for instructions or questions on how to use the provider portal or how to submit an ICE file:

[NetworkQuality@scanhealthplan.com](mailto:NetworkQuality@scanhealthplan.com)

### Rheumatoid Arthritis ICD-10 Codes

M05.0X – Felty's syndrome
M05.1X – Rheumatoid lung disease with rheumatoid arthritis
M05.2X – Rheumatoid vasculitis with rheumatoid arthritis
M05.3X – Rheumatoid heart disease with rheumatoid arthritis
M05.4X – Rheumatoid myopathy with rheumatoid arthritis
M05.5X – Rheumatoid polyneuropathy with rheumatoid arthritis
M05.6X – Rheumatoid arthritis with involvement of other organs and systems
M05.7X – Rheumatoid arthritis with rheumatoid factor of site without organ or systems involvement
M05.8X – Other rheumatoid arthritis with rheumatoid factor
M05.9 – Rheumatoid arthritis with rheumatoid factor
M06.0X – Rheumatoid arthritis without rheumatoid factor
M06.1 – Adult-onset Still's disease
M06.2X – Rheumatoid bursitis
M06.3X – Rheumatoid nodule
M06.8X – Other specified rheumatoid arthritis
M06.9 – Rheumatoid arthritis, unspecified

"X" indicates a placeholder to further specify etiology, anatomic site, severity and other code extensions.